



ATP2C1 gene

ATPase secretory pathway Ca²⁺ transporting 1

Normal Function

The *ATP2C1* gene provides instructions for making a protein called hSPCA1. This protein is an adenosine triphosphate (ATP)-powered calcium pump, which uses energy from ATP molecules to pump charged calcium atoms (calcium ions) across cell membranes. Specifically, the hSPCA1 protein transports calcium ions into a cell structure called the Golgi apparatus, where they are stored until needed. The appropriate storage and release of calcium is essential for many cell activities, including cell growth and division, cell movement, and attachment of cells to one another (cell adhesion).

The hSPCA1 protein also transports manganese ions into the Golgi apparatus. Manganese works with a variety of enzymes and is involved in processing newly formed proteins.

The hSPCA1 protein is present in cells throughout the body. It appears to be particularly important for the normal function of cells called keratinocytes, which are found in the outer layer of the skin (the epidermis).

Health Conditions Related to Genetic Changes

benign chronic pemphigus


More than 100 mutations in the *ATP2C1* gene have been found to cause benign chronic pemphigus, a rare skin condition characterized by red, raw, and blistered areas of skin. Mutations in this gene reduce the amount of functional hSPCA1 protein, which impairs the storage of calcium ions in the Golgi apparatus. For unknown reasons, this abnormal calcium storage affects keratinocytes more than other types of cells. Problems with calcium regulation impair many cell functions, including cell adhesion. As a result, keratinocytes do not stick tightly to one another, which causes the epidermis to become fragile and less resistant to minor trauma. Because the skin is easily damaged, it develops raw, blistered areas, particularly in skin folds where there is moisture and friction.

Although *ATP2C1* gene mutations probably also affect the transport of manganese within cells, abnormal manganese regulation is not thought to contribute to the signs and symptoms of benign chronic pemphigus.

Chromosomal Location

Cytogenetic Location: 3q22.1, which is the long (q) arm of chromosome 3 at position 22.1

Molecular Location: base pairs 130,850,500 to 131,016,712 on chromosome 3 (Homo sapiens Annotation Release 108, GRCh38.p7) (NCBI)



Credit: Genome Decoration Page/NCBI

Other Names for This Gene

- AT2C1_HUMAN
- ATP-dependent Ca(2+) pump PMR1
- ATP2C1A
- ATPase 2C1
- ATPase, Ca(2+)-sequestering
- ATPase, Ca++ transporting, type 2C, member 1
- BCPM
- calcium-transporting ATPase type 2C member 1
- HHD
- hSPCA1
- HUSSY-28
- KIAA1347
- PMR1
- secretory pathway Ca2+/Mn2+ ATPase 1
- SPCA1

Additional Information & Resources

Educational Resources

- Madame Curie Bioscience Database: The Golgi Apparatus
<https://www.ncbi.nlm.nih.gov/books/NBK6268/>

Scientific Articles on PubMed

- PubMed
<https://www.ncbi.nlm.nih.gov/pubmed?term=%28ATP2C1%5BTIAB%5D%29+AND+%28%28Genes%5BMH%5D%29+OR+%28Genetic+Phenomena%5BMH%5D%29%29+AND+english%5BIa%5D+AND+human%5Bmh%5D+AND+%22last+3600+days%22%5Bdp%5D>

OMIM

- ATPase, Ca(2+)-TRANSPORTING, TYPE 2C, MEMBER 1
<http://omim.org/entry/604384>

Research Resources

- ClinVar
<https://www.ncbi.nlm.nih.gov/clinvar?term=ATP2C1%5Bgene%5D>
- HGNC Gene Family: ATPases Ca²⁺ transporting
<http://www.genenames.org/cgi-bin/genefamilies/set/1209>
- HGNC Gene Symbol Report
http://www.genenames.org/cgi-bin/gene_symbol_report?q=data/hgnc_data.php&hgnc_id=13211
- NCBI Gene
<https://www.ncbi.nlm.nih.gov/gene/27032>
- UniProt
<http://www.uniprot.org/uniprot/P98194>

Sources for This Summary

- OMIM: ATPase, Ca(2+)-TRANSPORTING, TYPE 2C, MEMBER 1
<http://omim.org/entry/604384>
- Behne MJ, Tu CL, Aronchik I, Epstein E, Bench G, Bikle DD, Pozzan T, Mauro TM. Human keratinocyte ATP2C1 localizes to the Golgi and controls Golgi Ca²⁺ stores. J Invest Dermatol. 2003 Oct;121(4):688-94.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/14632183>
- Dobson-Stone C, Fairclough R, Dunne E, Brown J, Dissanayake M, Munro CS, Strachan T, Burge S, Sudbrak R, Monaco AP, Hovnanian A. Hailey-Hailey disease: molecular and clinical characterization of novel mutations in the ATP2C1 gene. J Invest Dermatol. 2002 Feb;118(2):338-43.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/11841554>

- Hu Z, Bonifas JM, Beech J, Bench G, Shigihara T, Ogawa H, Ikeda S, Mauro T, Epstein EH Jr. Mutations in ATP2C1, encoding a calcium pump, cause Hailey-Hailey disease. *Nat Genet.* 2000 Jan;24(1):61-5.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/10615129>
- Missiaen L, Dode L, Vanoevelen J, Raeymaekers L, Wuytack F. Calcium in the Golgi apparatus. *Cell Calcium.* 2007 May;41(5):405-16. Epub 2006 Nov 30. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/17140658>
- Missiaen L, Raeymaekers L, Dode L, Vanoevelen J, Van Baelen K, Parys JB, Callewaert G, De Smedt H, Segaert S, Wuytack F. SPCA1 pumps and Hailey-Hailey disease. *Biochem Biophys Res Commun.* 2004 Oct 1;322(4):1204-13. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/15336968>
- Sudbrak R, Brown J, Dobson-Stone C, Carter S, Ramser J, White J, Healy E, Dissanayake M, Larrègue M, Perrussel M, Lehrach H, Munro CS, Strachan T, Burge S, Hovnanian A, Monaco AP. Hailey-Hailey disease is caused by mutations in ATP2C1 encoding a novel Ca(2+) pump. *Hum Mol Genet.* 2000 Apr 12;9(7):1131-40.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/10767338>

Reprinted from Genetics Home Reference:
<https://ghr.nlm.nih.gov/gene/ATP2C1>

Reviewed: May 2013

Published: March 21, 2017

Lister Hill National Center for Biomedical Communications
U.S. National Library of Medicine
National Institutes of Health
Department of Health & Human Services